

DOCUMENT SCAN CONTROL SHEET

Document ID 509040

EPA # 1001-70

EPA #
For SLN

Product 3336 G
Name

Accepted
Date 11/10/2006

Document
Type PRILAB

Registration ID 115701

ACCEPTED
FOR REGISTRATION
NOV 10 2006
NYSDEC

509040

3336 G

TUR
ORN
SYS-

3336 G

TURF AND ORNAMENTAL SYSTEMIC FUNGICIDE

For the prevention and control of turf diseases and the diseases of annual and perennial flowers, bedding plants, foliage plants, ground covers, plus deciduous and evergreen trees and shrubs.

ACTIVE INGREDIENT:

Thiophanate-methyl (dimethyl 4,4'-o-phenylenebis(3-thioallophanate))

OTHER INGREDIENTS 2.08%
..... 97.92%

TOTAL 100.00%

VERSION 9.20.06

AMENDED 9.20.06

REPLACES 12/99

EPA REG. NO. 1001-70

EPA EST. NO. 961-PA-1

FIRST AID

- IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.
- IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for further treatment advice.
- IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes; then continue rinsing eye. Call a poison control center or doctor for further treatment advice.
- IF SWALLOWED: Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
- Have the product container or label with you when calling a poison control center or doctor, or when going for treatment.
For 24-Hour Emergency Assistance Call Chemtrec 1-800-424-9300.



CLEARY CHEMICAL CORPORATION
178 RIDGE ROAD
DAYTON, NJ 08810-1501
(732) 329-8399 (800) 524-1662

Keep Out Of Reach Of Children

CAUTION

URF AND

ORNAMENTAL

CLASSIFIED FOR
"RESTRICTED USE"
IN NEW YORK STATE
UNDER 6NYCRR PART 326

RF AND NAMENTAL STEMIC FUNGICIDE

CAUTION

Container Disposal: Shake empty product container to remove all possible material. When bags are empty, they may be disposed of in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

COMMERCIAL TURF AND ORNAMENTAL

GENERAL INFORMATION

3336 G is a broad spectrum fungicide exhibiting preventative, curative and systemic properties. It is useful on a wide variety of turf and ornamental disease problems. Apply 3336 G with ground based (hand operated/push) or mechanical (power) equipment sufficient to accurately deliver this product to the application site. Do not apply by hand or with hand held rotary spreader (e.g. belly grinder). Application equipment should be properly calibrated prior to use. Application equipment should be thoroughly cleaned after use. If a rate range is given, use the higher rate under conditions of severe disease pressure. Contact your local State Extension Service for recommendations for application schedules appropriate to your area. This product is not for homeowner use and can only be used by individuals/firms licensed or registered by the state to apply ornamental or turf pest control products. No claim of compatibility with other pesticides is implied. Do not apply this product through any type of irrigation system.

Resistance Management: To avoid the development of tolerant strains of fungi, 3336 G should be used with fungicides of different modes of action. Clearly, chemical does not recommend the use of products containing thiabendazole in combination or rotation with 3336 G. These utilize similar chemistry and mode of action and can contribute to development of disease tolerance. If, after using 3336 G as recommended, the treatment is not effective, a tolerant strain of fungi may be present. Consult your local Cleary Chemical representative, your State Agricultural Experiment Station, or your State Cooperative Extension Service for proper disease identification and advice on the prompt use of some other suitable fungicide or disease control strategy. As long as recommended precautions are followed, 3336 G can remain useful for disease control.

COMMERCIAL TURF APPLICATIONS

For use on all fine turf applications such as commercial, residential and public (such as home lawns, parks, athletic fields, schools, and day care centers), and golf courses (greens, tees, fairways and aprons) of cool and warm season grasses such as Bentgrass, Bluegrass, Bermudagrass, Fescue, Myegrass, St. Augustinegrass, Zoysiagrass, or their mixtures. 3336 G is not phytotoxic to any of the above mentioned grasses when used in accordance with the label. 3336 G is to be used for the prevention and control of the diseases mentioned below. 3336 G may be applied to dry or moist leaf/soil conditions and should be activated with irrigation or natural rainfall within 24 hours for best results. If irrigated, application should be watered in by applicator with 0.1-0.25 inches of water to begin activation of product for disease control. When treating golf greens, treatment of aprons is recommended. Use the highest recommended rate under conditions of severe disease pressure. Do not graze animals on treated turf. Do not feed clippings to livestock or poultry. Not for use on turf being grown for sale or other commercial use as sod.

Table 1: Maximum Individual Application Rates and Minimum Re-Treatment Intervals
Do not exceed the amounts per acre or reduce the re-treatment interval indicated below.

Use Site	Maximum application rate of 3336 G	Minimum Re-Treatment Interval	Comments
Residential or Public Areas	131 Lbs/Acre (3 lbs / 1,000 sq.ft.)	14 days	
Golf Course Tees Greens, Aprons	392 Lbs/Acre (9 lbs / 1,000 sq.ft.)	14 days	
Golf Course Fairways-- except Florida	262 Lbs/Acre (6 lbs / 1,000 sq.ft.)	14 days	Excludes Florida
Golf Course Fairways-- Florida Only	131 Lbs/Acre (3 lbs / 1,000 sq.ft.)	14 days	Florida Only One Application per year.

Table 2: Maximum Seasonal Application Rates
Do not exceed the following amounts of product per acre per season

Use Site	Maximum Pounds 3336 G per Acre per Season	Pounds 3336 G per 1,000 sq.ft.	Comments
Residential or Public Areas	523 lbs.	12 lbs.	4 Applications per year.
Golf Course Tees Greens, Aprons	1048 lbs.	24 lbs.	
Golf Course Fairways-- except Florida	262 lbs.	6 lbs.	Excludes Florida
Golf Course Fairways-- Florida Only	131 lbs.	3 lbs.	Florida Only

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed, absorbed through the skin or inhaled. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing dust or spray mist. Wash thoroughly with soap and water after handling.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

WPS Uses: Applicators and other handlers who handle this pesticide for any use covered by the Worker Protection Standard, 40 CFR Part 170, must wear long-sleeved shirt, long pants, and shoes with socks.
NON-WPS Uses: Applicators and other handlers who handle this pesticide for any use NOT covered by the Worker Protection Standard, 40 CFR Part 170, should wear appropriate clothing to reduce prolonged situations indicated in precautionary statements such as: long-sleeved shirt, long pants, and shoes with socks.
Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water by disposal of equipment washwaters. Do not apply, allow to drift, or drain or flush equipment onto non-target areas.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any other requirements specific to your State or Tribe, consult the agency responsible for pesticides.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific statements and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval (REI). The requirements in this box apply to uses of this product that are covered by the Worker Protection Standard.

Do Not enter or allow entry into treated areas during the restricted entry interval (REI) of 12 hours.

Personal Protective Equipment (PPE) required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water: is coveralls, chemical resistant gloves made of any waterproof material, and shoes with socks.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box only apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Turf and Landscape Uses: Do not enter or allow entry into treated areas until granules and dust have settled unless wearing personal protective equipment listed on this label.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in the original container in a dry, temperature controlled area. Do not store in a manner where cross-contamination with other pesticides, fertilizers, food or feed could occur. If spilled during storage or handling, contain/re-capture spillage and use in accordance with instruction below or dispose of in accordance with the Pesticide Disposal Instructions listed below.

Pesticide Disposal: Wastes resulting from the use of this product are toxic and may be disposed of on site or at an approved waste disposal facility.

360[®] B60

PERMANENTAL SYSTEMIC FUNGICIDE

the diseases of annual and perennial flowers, bedding plants, ground covers, plus deciduous

FIRST AID

HALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for further treatment advice.

EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for further treatment advice.

ALLOWED: Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

the product container or label with you when calling a poison control center or doctor, when going for treatment.

Hour Emergency Assistance Call Chemtrec 1-800-424-9300.



CLEARY CHEMICAL CORPORATION
178 RIDGE ROAD
DAYTON, NJ 08810-1501
(732) 329-8339 (800) 524-1662

Out Of Reach Of Children

Table 3: Turf Disease Control

Disease(s) Controlled	Rate of 3336 G lbs/1000 sq.ft.	Remarks*
Anthracnose, basal <i>Colletotrichum graminicola</i>	6-9	For prevention in historic areas of disease pressure, apply twice at 14 day intervals when soil temperature reaches 60°F. For curative control, apply when disease first appears in spring. Combine all of the above fungicides with chlorothalol or triadimefon can be utilized.
Anthracnose, foliar <i>Colletotrichum graminicola</i>	3-6	
Bermudagrass Decline <i>Guenamanomyces graminis</i> var. <i>graminis</i> <i>Guenamanomyces graminis</i> var. <i>avenae</i>	6-9	Apply in mid-July or when disease symptoms first appear and repeat at 14 day intervals for suppression. Use higher rates under most severe disease conditions to maintain plant vigor.
Cool Season Brown Patch <i>Rhizoctonia cerealis</i> Necrotic Ring Spot <i>Rhizoctonia cerealis</i> Spring Dead Spot <i>Leptospheria korrae</i>	6-9	For prevention, apply in fall before turf has stopped all growth activity. Apply second application in early spring when soil temperatures reach 55-60°F. For curative control, apply when disease first appears in early spring and continue at 14 day intervals.
Coprinus Snow Mold <i>Coprinus psychromorbidus</i>	6-9	Apply 2 treatments at 21 day intervals in late fall to early winter, with the last application made just before the winter snow cover. Rotations with PCNB can be utilized.
Dollar spot <i>Moellerodiscus</i> <i>Lonizia</i> , <i>Sclerotinia</i> Large Brown Patch <i>Dothidea</i> spp. Ascochyta Leaf Blight <i>Ascochyta</i> spp. Copper Spot <i>Glaesporangium sargisi</i> Large Brown Patch <i>Dothidea</i> spp. Fusarium Patch <i>Fusarium nivale</i> <i>Fusarium</i> spp. <i>Rhizoctonia solani</i>	1.5-6	For prevention, use lowest rate and apply application before expected period of disease development. Apply when disease first appears and continue at 14 day intervals. Rotations with chlorothalol, iprodione, or mancozeb (Protect™) can be utilized.
Fusarium Blight <i>Fusarium roseum</i> and <i>F. triticum</i>	6-9	Apply when disease first appears at 14 day intervals.
Gray Leaf Spot (Blast) <i>Pyricularia grisea</i>	6-9	Apply preventative application before expected period of disease development. Continue applications at 14 day intervals as needed.
Leaf Spot <i>Dothidea</i> spp. Leaf, crown and root diseases <i>Bipolaris</i> , <i>Cirularia</i> , and <i>Exserohilum</i> Pink Snow Mold <i>Microdochium nivale</i>	6-9	Apply when disease first appears and make applications at 14 day intervals. For prevention, apply in spring and fall. For prevention, apply 3 applications, starting first in early spring, at 14 day intervals. Rotations with chlorothalol, iprodione, or mancozeb (Protect) are recommended under severe conditions.
Stripes Smut <i>Ustilago striiformis</i>	6-9	Apply middle-late fall to early winter before turf has stopped all growth activity or in spring during periods of crown or disease symptoms. For curative control, apply at 14 day intervals. Best results are achieved when first disease period. Best results are achieved when first application is made before turf is completely dormant.
Summer Patch <i>Magnaporthe poae</i>	6-9	Apply when disease first appears and make two applications at 14 day intervals. For prevention, apply in spring and fall. For prevention, apply 3 applications, starting first in early spring, at 14 day intervals. Rotations with chlorothalol, iprodione, or mancozeb (Protect) may be used as part of the three application program. For suppression, apply at 14 day intervals when disease first appears.
Bentgrass Dead Spot <i>Ophiopogon distachyoides</i>	6-9	For prevention, apply in early June or based upon local conditions. For curative control, apply at 14 day intervals as needed. Rotations and/or tank mix combinations may be used for season long disease prevention.

*Observe the maximum individual application rates and maximum seasonal application limits in Table 2.

Table 4: Guide to 3336 G Recommended Spreader Settings*

These settings are presented as a guide for use. Age, condition of spreader and operator speed will cause variation in applied rates of material. Spreader calibration at the beginning of each application is strongly recommended.

SPREADER	Pounds of 3336G Applied per 1,000 Square Feet								
	2	3	4	5	6	7	8	9	
Central Drop	7	8	9						
Cyclone Rotary	4	5	5.5	6.5	6.75	7	7.5	8	
Earthway Drop	10	11	12						
Earthway Rotary	16	17	18						
Ezee-Flow	C3	C4							
Frank's Drop	6								
Gandy	25	28							
Greenview Drop-Spread II	2.5	3	3.5	4	4.5	5			
Greenview Drop-Spread III	4	6	6.6	7.2	7.8	8.4	9		

PERMANENTAL SYSTEMIC FUNGICIDE

TURF CARE SYSTEMS

CAUTION

Container Disposal: Shake empty product container to remove all possible material. When bags are empty they may be disposed of in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

COMMERCIAL TURF AND ORNAMENTAL

GENERAL INFORMATION

3336 G is a broad spectrum fungicide exhibiting preventative, curative and systemic properties. It is useful on a wide variety of turf and ornamental disease problems. Apply 3336 G with ground based hand operated (push) or mechanical (powered) equipment sufficient to accurately deliver this product to the application site. Do not apply by hand or with hand held rotary sprayer (e.g. belly grinder). Application equipment should be properly calibrated prior to use. Application equipment should be thoroughly cleaned after use. If rate angle is given, use the higher rate under conditions of severe disease pressure. Contact your local State Extension Service for recommendations for application schedules appropriate to your area. This product is not for homeowner use and can only be used by individuals/firms licensed or registered by the state to apply ornamental or turf pest control products. No claim of compatibility with other pesticides is implied. Do not apply this product through any type of irrigation system.

Resistance Management: To avoid the development of tolerant strains of fungi, 3336 G should be used with fungicides of different modes of action. Cleary Chemical does not recommend the use of products containing thiabendazole in combination or rotation with 3336 G. These utilize similar chemistry and mode of action and can contribute to development of disease tolerance. If, after using 3336 G as recommended, the treatment is not effective, a tolerant strain of fungi may be present. Consult your local Cleary Chemical representative, your State Agricultural Experiment Station, or your State Cooperative Extension Service for proper disease identification and advice on the prompt use of some other suitable fungicide or disease control strategy. As long as recommended precautions are followed, 3336 G can remain useful for disease control.

COMMERCIAL TURF APPLICATIONS

For use on all fine turf applications such as commercial, residential and public (such as home lawns, parks, athletic fields, schools, and day care centers), and golf courses (greens, tees, fairways and aprons) for cool and warm season grasses such as Bentgrass, Bluegrass, Bermudagrass, Rescue, Ryegrass, St. Augustinegrass, Zoysiagrass, or their mixtures. 3336 G is not phytotoxic to any of the above mentioned grasses when used in accordance with the label. 3336 G is to be used for the prevention and control of the diseases mentioned below. 3336 G may be applied to dry or moist leaf/soil conditions and should be activated with irrigation or natural rainfall within 24 hours for best results. If irrigated, application should be watered in by applicator with 0.1-0.25 inches of water to begin activation of product for disease control. When treating golf greens, treatment of aprons is recommended. Use the highest recommended rate under conditions of severe disease pressure. Do not graze animals on treated turf. Do not feed clippings to livestock or poultry. Not for use on turf being grown for sale or other commercial use as sod.

Table 1: Maximum Individual Application Rates and Minimum Re-Treatment Intervals
Do not exceed the amounts per acre or reduce the re-treatment interval indicated below.

Use Site	Maximum application rate of 3336 G	Minimum Re-Treatment Interval	Comments
Residential or Public Areas	131 Lbs/Acre (3 lbs / 1,000 sq.ft.)	14 days	
Golf Course Tees, Greens, Aprons	303 Lbs/Acre (9 lbs / 1,000 sq.ft.)	14 days	
Golf Course Fairways—except Florida	262 Lbs/Acre (6 lbs / 1,000 sq.ft.)	14 days	Excludes Florida
Golf Course Fairways—Florida Only	131 Lbs/Acre (3 lbs / 1,000 sq.ft.)	14 days	Florida Only One Application per year.

Table 2: Maximum Seasonal Application Rates
Do not exceed the following amounts of product per Acre per season

Use Site	Maximum Pounds 3336 G per Acre per Season	Pounds 3336 G per 1,000 sq.ft.	Comments
Residential or Public Areas	523 lbs.	12 lbs.	4 Applications per-year
Golf Course Tees, Greens, Aprons	1048 lbs.	24 lbs.	
Golf Course Fairways—except Florida	262 lbs.	6 lbs.	Excludes Florida
Golf Course Fairways—Florida Only	131 lbs.	3 lbs.	Florida Only

moderate eye irritation. Avoid contact with soap and water after handling.

use covered by the Worker Protection with socks. If any use NOT covered by the Worker use prolonged situations indicated in h socks.

tions for washables, use detergent

wing gum, using tobacco or using roughly and put on clean clothing. wash thoroughly and change into

urface water is present, or to intertidal equipment washwaters. Do not apply

ENT WITH ITS LABELING. Do not apply through drift. Only protected handlers ate or Tribe, consult the agency respon-

ction Standard, 40 CFR part 170. This is forests, nurseries, and greenhouses, amination, notification, and emergency ations on this label about personal this box apply to uses of this product

(RE) of 12 hours. permitted under the Worker Protection ents, soil, or water, is, coveralls, chemical

VTS in the scope of the Worker Protection product is used to produce agricultural

les and dust have settled unless wearing

a. Do not store in a manner where cross-storage or handling, contain/re-capture with the Pesticide Disposal Instructions

be disposed of on site or at an approved

Greenview Drop-Spread II	2.5	3	3.5	4	4.5	5
Greenview Jet-Spread II	4	5	5.5	6.5	7	8
Greenview Jet-Spread III	4.5	5.5	6	7	7.5	8
Greenview Snow 'N Tell	7	8	9	10		
Jackson Drop	5.5		6.5			
Lesco	G	1	J	K	L	M
Orbita Drop	5	7	8	9	10	10.5
Orbita Rotary	4	5	7	8	9	10
Precision Drop						
Red Devil Rotary	7					
Republic Drop	5	7	8	9	10	10.5
Republic Rotary	4	5	7	8	9	10
Scott's AccuGreen	6	6.5	7			
Scott's Drop PFI 1	5	6	7	8	9	9.5
Scott's Easy Green	2.5	2.6	2.7			
Scott's Easy Gro	7.5		8			
Scott's RX7 Rotary	1	J	K	L	M	N
Scott's RBA Rotary	K	L	M	N	O	P
Scott's Rotary Retail	5.5	6	6.5	6.5	7	7.5
Scott's Speedy Green	5.5	5.75	6	8.5		
Seas Craftsman Drop	7	7.5	8			
Spyker	2.5	3	3.5			

*SGN = 118

COMMERCIAL HORTICULTURAL APPLICATIONS

Nursery, Greenhouse, Landscape & Interiorscape Annual and Perennial Flowers, Bedding Plants, Foliage Plants, Ground Covers, plus Deciduous and Evergreen Trees and Shrubs

Do not apply to home orchards/fruit trees after fruit set.

3336 G is a broad spectrum systemic fungicide which controls a variety of foliar, stem and root diseases on a wide range of commercially important plants. 3336 G may be used for disease control and prevention programs on annual and perennial flowers, bedding plants, foliage plants, ground covers, and deciduous and evergreen trees and shrubs found in Nurseries, Greenhouses, Interiorscapes, and Landscape sites. Follow guidelines provided under general information for use of this product.

Application Instructions: 3336 G may be used as a broadcast, fly-on, surface, over the top, and media incorporation application for the diseases listed below. For surface and broadcast applications to established plants apply at the rates below; 3336 G may be applied to dry or moist soil conditions and should be activated with irrigation or natural rainfall within 24 hours for best results. If irrigated, applications should be watered in by applicator with 0.1-0.25 inches of water to begin activation. For media incorporation, mix at a rate of 2-3 lbs/100 yd before seeding or transplanting of desired crop. Media incorporated with 3336 G may be stockpiled for later use without activation until needed. For ground applications, do not exceed 300 pounds active ingredient per acre per crop season from all thiophanate-methyl containing products.

Note: The "Directions For Use" of this product reflect the cumulative inputs from both historical field use and product testing programs. However, it is impossible to test this product on all species and cultivars. A preliminary trial is suggested on a small scale before a full treatment is applied to any plant type not shown on this label but found in a similar use site with a listed disease problem. Wait 5-7 days after treatment to evaluate results. This product is not recommended for use on Swedish Ivy (*Nephtrolepis exaltata*), Boston Fern (*Polystichum australe*), and Easter Cactus (*Hatiora gaertneri*).

Diseases(s) Controlled

Rate of 3336 G lbs/1000 sq.ft.

For control of soil borne diseases of plants caused by: *Botrytis*, *Cyrtospora*, *Phytophthora*, *Pyrenopeziza*, *Rhizoctonia*, *Sclerotinia*, and *Thielaviopsis*.
For preventative control apply as media incorporation over the top application after seeding or transplanting to propagation beds, containers, pots, or nursery or landscape beds. For curative control, apply after seeding or transplanting when disease first appears. Note: 3336 G does not control *Pythium* or *Phytophthora* rotations with metaxyl, mefenoxam, etridiazole, propanil, mono and dipotassium phosphites. Do not use 3336 G for control of *Pythium* and *Phytophthora*.

3336[®] is a registered trademark of Cleary Chemical Corporation. Procept[™] is a trademark of Cleary Chemical Corporation. Copyright © 2006 Cleary Chemical Corporation.

3336 G

TURF ORNA- MENTALS

3336

TURF AND ORNAMENTALS

For the prevention and control of turf diseases and the diseases of annual and perennial flowers and evergreen trees and shrubs.

ACTIVE INGREDIENT:

Thiophanate-methyl (dimethyl 4,4'-o-phenylenebis(3-methylphosphonate))

2.08%

OTHER INGREDIENTS

97.92%

TOTAL

100.00%

VERSION: 9/20/06

AMENDED: 9/20/06

REPLACES: 12/99

EPA REG. NO. 1001-70

EPA EST. NO. 961-PA-1

FIRST AID

- IF INHALED: Move person to fresh air. If person is not breathing, give artificial respiration, preferably by mouth-to-mouth respiration. Call poison control center or doctor for further treatment advice.
- IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse with water for 15-20 minutes. Call a poison control center or doctor for further treatment advice.
- IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Call a poison control center or doctor for further treatment advice.
- IF SWALLOWED: Call poison control center or doctor immediately. Do not induce vomiting. If person is conscious, have person sip a glass of water if able to swallow. Do not induce vomiting. Call a poison control center or doctor. Do not give anything by mouth to an unconscious person.
- Have the product container or label with you when calling a poison control center or doctor for treatment information.

For 24-Hour Emergency Assistance Call Chemtrec: 1-800-424-9

Keep Out Of Reach



FAL FUNGICIDE

Remove PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

● ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water by disposal of equipment/washwaters. Do not apply, allow to drift, or drain or flush equipment onto non-target areas.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any other requirements specific to your State or Tribe, consult the agency responsible for pesticides.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific statements and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval (REI). The requirements in this box apply to uses of this product that are covered by the Worker Protection Standard.

Do Not enter or allow entry into treated areas during the restricted entry interval (REI) of 12 hours.

Personal Protective Equipment (PPE) required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is coveralls, chemical resistant gloves made of any waterproof material, and shoes with socks.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box only apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Turf and Landscape Uses: Do not enter or allow entry into treated areas until granules and dust have settled unless wearing personal protective equipment listed on this label.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in the original container in a dry, temperature controlled area. Do not store in a manner where cross-contamination with other pesticides, fertilizers, food or feed could occur. If spilled during storage or handling, contain/re-capture spillage and use in accordance with instruction below or dispose of in accordance with the Pesticide Disposal Instructions listed below.

Pesticide Disposal: Wastes resulting from the use of this product are toxic and may be disposed of on site or at an approved waste disposal facility.

LIMITED WARRANTY AND DISCLAIMER

CLEARLY CHEMICAL CORPORATION warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use, subject to the risks referred to therein. CLEARLY CHEMICAL CORPORATION does not warrant the fitness, quality, or quantity of the material. CLEARLY CHEMICAL CORPORATION does not warrant any implied warranty of fitness or merchantability or any other expressed or implied warranties. To the extent consistent with applicable law neither CLEARLY CHEMICAL nor seller shall be liable for consequential, special or indirect damages resulting from the use or handling of this product, including, but not limited to, loss of profits, business reputation, or customers, labor costs, or other expenses incurred in planting or harvesting. CLEARLY CHEMICAL and seller offer this product and the buyer and user accept it subject to the foregoing conditions of sale and warranty which may be voided only by agreement in writing signed by a duly authorized representative of CLEARLY CHEMICAL CORPORATION.